



Cosmetic & Reconstructive Periodontics
Dental Implants

Richard H. Yamada, D.D.S
Douglas V. Gorin, D.D.S
Richard F. Marinello, D.D.S
Mark A. Rosen, D.D.S
Stephen P. Russo, D.M.D

POST-OPERATIVE CARE

CARE OF THE MOUTH

Brushing and Flossing: Begin brushing and flossing the areas not operated on the day following surgery. AVOID THE SURGICAL SITES until given specific brushing instructions at your post-operative visit. You may notice a white film over the surgical sites. This is completely normal.

Mouthwash: Use the prescribed mouthwash twice daily beginning the morning after surgery until instructed to discontinue.

Electric Toothbrushes: DO NOT use an electric toothbrush such as Sonicare or Oral-B Braun for six weeks following surgery.

Water Pik: DO NOT use a Water Pik for at least six weeks following surgery.

EATING

First 24 hours: Adequate nutrition is essential for normal healing. Following surgery, your diet should be restricted to COLD LIQUIDS. Dietary supplements such as Instant Breakfast or Ensure are good choices. AVOID PEROXIDE, ALCOHOL, CARBONATED BEVERAGES, AND DRINKING THROUGH A STRAW.

After 24 hours: Gradually progress to soft foods such as cooked vegetables, baked fish, tuna, pasta and meatloaf, which are easily chewed and swallowed. DO NOT SKIP MEALS. If you take nourishment regularly, you will feel better, gain strength, have less discomfort, and heal faster.

ACTIVITIES

Rest: Plan to rest at home the remainder of the day of surgery and the following day. You may read, watch TV or work at your desk at home. When sleeping, elevate your head to decrease swelling.

After 24 hours: You may return to your regular schedule, but avoid strenuous activities such as heavy lifting, jogging, exercise programs, etc. for one week following surgery.

MEDICATIONS

Pain Medications: Take pain medication within one hour after treatment with milk, fruit juice, or a full glass of water. Never take pain medication on an empty stomach. This medication may be repeated every three to four hours as needed for discomfort. **DO NOT USE ASPIRIN** or products containing aspirin following surgery. Aspirin, Advil and Aleve can contribute to bleeding problems. Only non-aspirin pain reliever, Tylenol is acceptable.

Antibiotics: Take prescribed antibiotics as directed until all have been taken.

Steroid Dose-Pak: If prescribed, take as directed. If the appointment is late in the day, it is important to take all six tablets the day of the surgery.

SWELLING

Ice Packs: Some swelling usually occurs after periodontal surgery. To minimize this, you will be advised to use ice packs. Place the ice pack on the outside of your face, over the treated area, for 10 minutes, and then take it off for 10 minutes. Continue to apply ice packs as much as possible the first 24 hours after surgery.

BLEEDING

Slight bleeding or oozing is normal during the first 1-2 days following surgery. If excessive bleeding occurs, please call our office.

SMOKING

DO NOT smoke for three weeks after surgery. Smoking is detrimental to healing tissues and will affect the results of surgery. If you can stop smoking for three weeks post-operatively, you may as well quit all together.

SUTURES

Resorbable: The resorbable sutures used usually will be resorbed by the time you return for your post-operative visit.

Non-Resorbable: If non-resorbable sutures were placed, they will be removed at your post-operative visit.

It is our sincere desire that you are as comfortable as possible following surgery. If you should have any questions or problems, please call our office anytime: Chicago – 312-641-2572 or Skokie – 847-675-7555.

AFTER HOURS - PAGER NUMBERS

Dr. Yamada	312-740-4000
Dr. Gorin	312-740-4001
Dr. Marinello	312-740-4002
Dr. Rosen	847-217-4780
Dr. Russo	312-371-3654
Dr. Doan	312-972-2930